

Full-Length

ADD3 (Human) Recombinant Protein (P01)

Catalog # H00000120-P01

Size 50 ug

Specification

Product Description	Human ADD3 full-length ORF (AAH62559.1, 1 a.a. - 706 a.a.) recombinant protein with GST-tag at N-terminal.
Sequence	MSSDASQGVITPPPPSPMPHKERYFDRINENDPEYIRERNMSPDLRQDFNMMEQRKRVTTQILQSP AFREDLECLIQEQMKKGHNPTGLLALQQIADYIMANSFSGFSSPPLSLGMVTPINDLPGADTSSYV KGEKLTRCKLASLYRLVDLFGWAHLANTYISVRISKEQDHIIIPRGLSFSEATASNLVKVNIIGEVVDQ GSTNLKIDHTGFSPHAAIYSTRPDVKCVIHIHTLATAAVSSMKCGILPISQESLLLGDVAYYDYQGS EEQEERIQLQKVLGPSCVKLVLRNHGVVALGETLEEFHYIFNVQLACEIQVQALAGAGGVDNLH VLDFQKYKAFTYTVAASGGGGVNMGSHQKWKVGEIEFEGLMRTLDNLGYRTGYAYRHPLIREKPR HKSDVEIPATVTAFSFEDDTVPLSPLKYMAQRQQREKTRWLNSPNTYMKVNVPEESRNGETSPR TKITWMKAEDSSKVSOGTPIKIEDPNQFVPLNTNPNEVLEKRNKIREQNRYDLKTAGPQSQLLAGIV VDKPPSTMQFEDDDHGPPAPPNPFSLHTEGELEEYKRTIERKQQGLEDAEQELLSDASSVSQI QSQTQSPQNVPEKLEENHELFSKSFISMEVPVMVVNGKDDMHDVEDELAKRVSRLSTSTTIENIEI TKSPEKIEEVLSPGSPSKSPSKKKKKFRTPSFLKKNKKKEKVEA
Host	Wheat Germ (in vitro)
Theoretical MW (kDa)	104.06
Interspecies Antigen Sequence	Mouse (92); Rat (93)
Preparation Method	in vitro wheat germ expression system
Purification	Glutathione Sepharose 4 Fast Flow
Storage Buffer	50 mM Tris-HCl, 10 mM reduced Glutathione, pH=8.0 in the elution buffer.
Storage Instruction	Store at -80°C. Aliquot to avoid repeated freezing and thawing.
Note	Best use within three months from the date of receipt of this protein.

Applications

- Enzyme-linked Immunoabsorbent Assay

- Western Blot (Recombinant protein)
- Antibody Production
- Protein Array

Gene Info — ADD3

Entrez GeneID [120](#)

GeneBank Accession# [BC062559.1](#)

Protein Accession# [AAH62559.1](#)

Gene Name ADD3

Gene Alias ADDL

Gene Description adducin 3 (gamma)

Omim ID [601568](#)

Gene Ontology [Hyperlink](#)

Gene Summary

Adducins are heteromeric proteins composed of different subunits referred to as adducin alpha, beta and gamma. The three subunits are encoded by distinct genes and belong to a family of membrane skeletal proteins involved in the assembly of spectrin-actin network in erythrocytes and at sites of cell-cell contact in epithelial tissues. While adducins alpha and gamma are ubiquitously expressed, the expression of adducin beta is restricted to brain and hematopoietic tissues. Adducin, originally purified from human erythrocytes, was found to be a heterodimer of adducins alpha and beta. Polymorphisms resulting in amino acid substitutions in these two subunits have been associated with the regulation of blood pressure in an animal model of hypertension. Heterodimers consisting of alpha and gamma subunits have also been described. Structurally, each subunit is comprised of two distinct domains. The amino-terminal region is protease resistant and globular in shape, while the carboxy-terminal region is protease sensitive. The latter contains multiple phosphorylation sites for protein kinase C, the binding site for calmodulin, and is required for association with spectrin and actin. Alternatively spliced adducin gamma transcripts encoding different isoforms have been described. The functions of the different isoforms are not known. [provided by RefSeq

Other Designations OTTHUMP00000020463|OTTHUMP00000020464|adducin-like protein 70

Disease

- [Alzheimer Disease](#)
- [Cardiovascular Diseases](#)

- [Diabetes Mellitus](#)
- [Edema](#)
- [Endolymphatic Hydrops](#)
- [Genetic Predisposition to Disease](#)
- [Hypertension](#)
- [Meniere Disease](#)
- [Tobacco Use Disorder](#)