

FKBP4 monoclonal antibody, clone Hi52C (FITC)

Catalog # MAB11384

Size 100 ug

Applications

Application Data with Unconjugated Antibody

Immunohistochemistry



Immunohistochemical staining of human prostate tissue with FKBP4 monoclonal antibody, clone Hi52C (Cat # MAB11372) shows the strongest staining for FKBP4 is in ductal epithelial cells.

Specification

Product Description Mouse monoclonal antibody raised against synthetic peptide of FKBP4.

Immunogen A synthetic peptide corresponding to residues of human FKBP4.

Host Mouse

Reactivity Dog, Hamster, Human, Mouse, Rat

Form Liquid

Conjugation FITC

Purification Protein G purification

Isotype IgG

Recommend Usage Western Blot (1:2000)
Immunohistochemistry (1:250)
Immunoprecipitation (5 ug with 10-20 uL Protein A beads)
The optimal working dilution should be determined by the end user.

Storage Buffer In PBS, pH 7.4 (50% glycerol, 0.09% sodium azide)

Storage Instruction Store at 4°C.
Aliquot to avoid repeated freezing and thawing.

Note

Application Data with Unconjugated Antibody
Immunohistochemistry
Immunohistochemical staining of human prostate tissue with FKBP4 monoclonal antibody, clone Hi5 2C (Cat # MAB11372) shows the strongest staining for FKBP4 is in ductal epithelial cells.

Applications

- Western Blot
- Immunohistochemistry
- Immunoprecipitation

Gene Info — FKBP4

Entrez GeneID [2288](#)

Protein Accession# [NP_002005.1](#)

Gene Name FKBP4

Gene Alias FKBP52, FKBP59, HBI, Hsp56, PPlase, p52

Gene Description FK506 binding protein 4, 59kDa

Omim ID [600611](#)

Gene Ontology [Hyperlink](#)

Gene Summary

The protein encoded by this gene is a member of the immunophilin protein family, which play a role in immunoregulation and basic cellular processes involving protein folding and trafficking. This encoded protein is a cis-trans prolyl isomerase that binds to the immunosuppressants FK506 and rapamycin. It has high structural and functional similarity to FK506-binding protein 1A (FKBP1A), but unlike FKBP1A, this protein does not have immunosuppressant activity when complexed with FK506. It interacts with interferon regulatory factor-4 and plays an important role in immunoregulatory gene expression in B and T lymphocytes. This encoded protein is known to associate with phytanoyl-CoA alpha-hydroxylase. It can also associate with two heat shock proteins (hsp90 and hsp70) and thus may play a role in the intracellular trafficking of hetero-oligomeric forms of the steroid hormone receptors. This protein correlates strongly with adeno-associated virus type 2 vectors (AAV) resulting in a significant increase in AAV-mediated transgene expression in human cell lines. Thus this encoded protein is thought to have important implications for the optimal use of AAV vectors in human gene therapy. The human genome contains several non-transcribed pseudogenes similar to this gene. [provided by RefSeq]

Other Designations

52 kD FK506 binding protein|FK506 binding protein 4 (59kD)|FK506 binding protein 52|FK506-binding protein 4 (59kD)|HSP binding immunophilin|T-cell FK506-binding protein, 59kD|p59 protein|peptidylprolyl cis-trans isomerase|rotamase

Disease

- [Asthma](#)
- [Genetic Predisposition to Disease](#)
- [Glaucoma](#)
- [Hypospadias](#)
- [Syndrome](#)