


RecomAb™

Mouse IgG monoclonal antibody, clone RM104 (Biotin)

Catalog # MAB14934 Size 50 ug

Applications




Western Blot

Western blot analysis of nonreduced (-) and reduced (+) mouse immunoglobulins (20 ng/lane) with Mouse IgG monoclonal antibody, clone RM104 (Biotin) (Cat # MAB14934) at 0.2 ug/mL working concentration. This antibody reacts to nonreduced mouse IgG1, IgG2a, IgG2b, and IgG3. It showed no cross reactivity with IgM, IgA, or IgE.

Enzyme-linked Immunoabsorbent Assay

A titer ELISA of mouse IgG. The plate was coated with different amounts of mouse IgG. A serial dilution of Mouse IgG monoclonal antibody, clone RM104 (Biotin) (Cat# MAB14934)4 was used as the primary antibody. An alkaline phosphatase conjugated anti-rabbit IgG as the secondary antibody.



Enzyme-linked Immunoabsorbent Assay

ELISA analysis of Mouse IgG monoclonal antibody, clone RM104 (Cat # MAB12765) at the following concentrations: 0.2 ug/mL, 0.05 ug/mL, and 0.01 ug/mL. The plate was coated with 50 ng/well of different immunoglobulins and the result shows this antibody reacts to the Fc region of mouse IgG1, IgG2a, IgG2b, and IgG3; no cross reactivity with IgM, IgA, IgE, human or rat IgG. An alkaline phosphatase conjugated anti-rabbit IgG as the secondary antibody.

Specification

Product Description

Rabbit recombinant monoclonal antibody raised against mouse IgG.

🖗 Abnova

Product Information

Antibody Species	Rabbit
Immunogen	Original antibody is raised against mouse IgG.
Sequence	N/A
Reactivity	Mouse
Form	Liquid
Conjugation	Biotin
Purification	Protein A purification
lsotype	lgG
Recommend Usage	ELISA (0.005-0.2 ug/mL) Flow Cytometry Immunocytochemistry (0.5-2 ug/mL) Immunohistochemistry (0.5-2 ug/mL) Immunoprecipitation Western Blotting (0.1-0.5 ug/mL) The optimal working dilution should be determined by the end user.
Storage Buffer	In PBS (50% glycerol, 1% BSA, 0.09% sodium azide)
Storage Instruction	Store at -20°C. Aliquot to avoid repeated freezing and thawing.
Note	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which shoul d be handled by trained staff only.

Applications

Western Blot

Western blot analysis of nonreduced (-) and reduced (+) mouse immunoglobulins (20 ng/lane) with Mouse IgG monoclonal antibody, clone RM104 (Biotin) (Cat # MAB14934) at 0.2 ug/mL working concentration. This antibody reacts to nonreduced mouse IgG1, IgG2a, IgG2b, and IgG3. It showed no cross reactivity with IgM, IgA, or IgE.

- Immunohistochemistry
- Immunocytochemistry
- Immunoprecipitation

Enzyme-linked Immunoabsorbent Assay

A titer ELISA of mouse IgG. The plate was coated with different amounts of mouse IgG. A serial dilution of Mouse IgG monoclonal antibody, clone RM104 (Biotin) (Cat# MAB14934)4 was used as the primary antibody. An alkaline phosphatase conjugated anti-rabbit IgG as the secondary antibody.

Enzyme-linked Immunoabsorbent Assay

ELISA analysis of Mouse IgG monoclonal antibody, clone RM104 (Cat # MAB12765) at the following concentrations: 0.2 ug/mL, 0.05 ug/mL, and 0.01 ug/mL. The plate was coated with 50 ng/well of different immunoglobulins and the result shows this antibody reacts to the Fc region of mouse IgG1, IgG2a, IgG2b, and IgG3; no cross reactivity with IgM, IgA, IgE, human or rat IgG. An alkaline phosphatase conjugated anti-rabbit IgG as the secondary antibody.

Flow Cytometry

Gene Info — Ighg1	
Entrez GenelD	<u>16017</u>
Protein Accession#	<u>P01868</u>
Gene Name	lghg1
Gene Alias	lgG1, lgh-4, VH7183
Gene Description	immunoglobulin heavy constant gamma 1 (G1m marker)
Gene Ontology	Hyperlink
Other Designations	-