

Goat Anti-Cat IgG F(ab')₂ secondary antibody (Rhodamine)

Catalog # PAB10469

Size 2 mg

Specification

Product Description	Goat anti-cat IgG recognizes cat IgG F(ab') ₂ fragment. This secondary antibody was purified using a ntigen affinity chromatography. The antibody is conjugated with Rhodamine.
Immunogen	Cat IgG F(ab') ₂ fragment
Host	Goat
Reactivity	Cat
Form	Lyophilized
Conjugation	Rhodamine (TRITC)
Purification	This product was prepared from monospecific antiserum by immunoaffinity chromatography, followed by solid phase adsorption(s) to remove any unwanted reactivities.
Recommend Usage	The optimal working dilution should be determined by the end user.
Storage Buffer	Lyophilized from 20 mM KH ₂ PO ₄ , 150 mM NaCl, pH 7.2
Storage Instruction	Store at 4°C on dry atmosphere. After reconstitution with 1 mL of deionized water, store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Applications

- Immunofluorescence
- Flow Cytometry

Publication Reference

- [Texas Red, a hydrophilic, red-emitting fluorophore for use with fluorescein in dual parameter flow microfluorometric and fluorescence microscopic studies.](#)

J A Titus, R Haugland, S O Sharrow, D M Segal.

Journal of Immunological Methods 1982 Apr; 50(2):193.