

# Rabbit Anti-Goat IgG (H&L) secondary antibody

Catalog # PAB9216

Size 500 uL

## Specification

|                            |   |
|----------------------------|---|
| <b>Product Description</b> | Rabbit anti-goat IgG recognizes goat IgG whole molecule. This secondary antibody was purified using antigen affinity chromatography.  |
| <b>Immunogen</b>           | Goat IgG whole molecule   |
| <b>Host</b>                | Rabbit  |
| <b>Reactivity</b>          | Goat  |
| <b>Specificity</b>         | No reaction was observed against anti-Pepsin or anti-rabbit IgG F(c).   |
| <b>Form</b>                | Lyophilized   |
| <b>Conjugation</b>         | Rhodamine (TRITC)   |
| <b>Purification</b>        | This product was prepared from monospecific antiserum by immunoaffinity chromatography, followed by solid phase adsorption(s) to remove any unwanted reactivities, pepsin digestion and chromatographic separation. |
| <b>Conjugation Note</b>    | 1.6 moles TRITC per mole of rabbit IgG F(ab') <sub>2</sub>  |
| <b>Recommend Usage</b>     | The optimal working dilution should be determined by the end user.  |
| <b>Storage Buffer</b>      | Lyophilized from 20 mM KH <sub>2</sub> PO <sub>4</sub> , 150 mM NaCl, pH 7.2 (0.01% sodium azide)   |
| <b>Storage Instruction</b> | Store at 4°C on dry atmosphere.<br>After reconstitution with 0.5 mL of deionized water, store at -20°C or lower.<br>Aliquot to avoid repeated freezing and thawing.   |
| <b>Note</b>                | This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.  |

## Applications

- Immunofluorescence