PGR polyclonal antibody

Catalog # PAB30378 Size 100 uL

Applications



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical staining (Formalin-fixed paraffin-embedded sections) of human uterine corpus with PGR polyclonal antibody (Cat # PAB30378) shows strong nuclear positivity in glandular cells at 1:200-1:500 dilution.

Specification

Product Description	Rabbit polyclonal antibody raised against partial recombinant human PGR.
Immunogen	Recombinant protein corresponding to human PGR.
Sequence	RVVRALDAVALPQPVGVPNESQALSQRFTFSPGQDIQLIPPLINLLMSIEPDVIYAGHDNTKPDTSS SLLTSLNQLGERQLLSVVKWSKS
Host	Rabbit
Reactivity	Human
Form	Liquid
Purification	Antigen affinity purification
Isotype	IgG
Recommend Usage	Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) (1:200-1:500) The optimal working dilution should be determined by the end user.
Storage Buffer	In PBS, pH 7.2 (40% glycerol, 0.02% sodium azide).

Copyright © 2023 Abnova Corporation. All Rights Reserved.

Storage Instruction	Store at 4°C. For long term storage store at -20°C. Aliquot to avoid repeated freezing and thawing.
Note	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which shoul d be handled by trained staff only.

Applications

- Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

  Immunohistochemical staining (Formalin-fixed paraffin-embedded sections) of human uterine corpus with PGR polyclonal antibody (Cat # PAB30378) shows strong nuclear positivity in glandular cells at 1:200-1:500 dilution.

Gene Info — PGR

Entrez GeneID	5241
Protein Accession#	P06401
Gene Name	PGR
Gene Alias	NR3C3, PR
Gene Description	progesterone receptor
Omim ID	607311
Gene Ontology	Hyperlink
Gene Summary	This gene encodes a member of the steroid receptor superfamily. The encoded protein mediates the physiological effects of progesterone, which plays a central role in reproductive events associ ated with the establishment and maintenance of pregnancy. This gene uses two distinct promotor s and translation start sites in the first exon to produce two isoforms, A and B. The two isoforms ar e identical except for the additional 165 amino acids found in the N-terminus of isoform A only, an d mediate their own response genes and physiologic effects with little overlap. The location of tran scription initiation for isoform B has not been clearly determined. [provided by RefSeq
Other Designations	-

Disease

- Abortion
- Adenocarcinoma

Copyright © 2023 Abnova Corporation. All Rights Reserved.

- Adhesions
- Aortic Aneurysm
- Arthritis
- Breast cancer
- Breast Neoplasms
- Carcinoma
- Cardiovascular Diseases
- Colorectal Neoplasms
- Connective Tissue Diseases
- Diabetes Mellitus
- Ductus Arteriosus
- Edema
- Endometrial Neoplasms
- Endometriosis
- Esophageal Neoplasms
- Fetal Diseases
- Gender Identity
- Genetic Predisposition to Disease
- Hyperprolactinemia
- Hypertension
- Infant
- Infection
- Infertility
- Inflammation
- Kidney Failure
- Leiomyoma

Copyright © 2023 Abnova Corporation. All Rights Reserved.

- Lung Neoplasms
- Lymphatic Metastasis
- Lymphoma
- Migraine Disorders
- Musculoskeletal Diseases
- Neoplasm Invasiveness
- Neoplasms
- Obesity
- Osteoporosis
- Ovarian cancer
- Ovarian Cysts
- Ovarian Failure
- Ovarian Neoplasms
- Panic Disorder
- Peritoneal Diseases
- Polycystic Ovary Syndrome
- Pregnancy Complications
- Premature Birth
- Prostate cancer
- Prostatic Neoplasms
- Proteinuria
- Psychiatric Status Rating Scales
- Puberty
- Pulmonary Disease
- Skin Diseases

Copyright © 2023 Abnova Corporation. All Rights Reserved.

- Stomach Neoplasms
- Thrombophilia
- Tobacco Use Disorder
- Urinary Bladder Neoplasms
- Uterine Diseases
- Uterine Neoplasms
- Vertigo
- Werner syndrome

Copyright © 2023 Abnova Corporation. All Rights Reserved.