RecomAb™

CXCL1 recombinant monoclonal antibody, clone R08-8D0

Catalog # RAB04406 Size 100 uL

Specification

Product Description	Rabbit recombinant monoclonal antibody raised against human CXCL1.
Antibody Species	Rabbit
Immunogen	Original antibody is raised against recombinant protein corresponding to human CXCL1.
Reactivity	Human
Form	Liquid
Preparation Method	Mammalian cell (CHO) expression system
Purification	Affinity purification
Isotype	IgG
Recommend Usage	ELISA (1:10000) The optimal working dilution should be determined by the end user.
Storage Buffer	In 50 mM Tris-Glycine pH 7.4 (0.15 M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% BSA).
Storage Instruction	Store at 4°C for at least six months Aliquot to avoid repeated freezing and thawing.

Applications

- Enzyme-linked Immunoabsorbent Assay

Gene Info — CXCL1

Entrez GeneID	2919

Copyright © 2023 Abnova Corporation. All Rights Reserved.

Protein Accession#	P09341
Gene Name	CXCL1
Gene Alias	FSP, GRO1, GROa, MGSA, MGSA-a, NAP-3, SCYB1
Gene Description	chemokine (C-X-C motif) ligand 1 (melanoma growth stimulating activity, alpha)
Omim ID	155730
Gene Ontology	Hyperlink
Gene Summary	Chemokines are a group of small (approximately 8 to 14 kD), mostly basic, structurally related molecules that regulate cell trafficking of various types of leukocytes through interactions with a subset of 7-transmembrane, G protein-coupled receptors. Chemokines also play fundamental roles in the development, homeostasis, and function of the immune system, and they have effects on cells of the central nervous system as well as on endothelial cells involved in angiogenesis or angiostasis. Chemokines are divided into 2 major subfamilies, CXC and CC, based on the arrangement of the first 2 of the 4 conserved cysteine residues; the 2 cysteines are separated by a single amino acid in CXC chemokines and are adjacent in CC chemokines. CXC chemokines are further subdivided into ELR and non-ELR types based on the presence or absence of a glu-leu-arg sequence adjacent and N terminal to the CXC motif. ELR types are chemotactic for neutrophils, while non-ELR types are chemotactic for lymphocytes.[supplied by OMIM
Other Designations	GRO1 oncogene (melanoma growth stimulating activity, alpha)|GRO1 oncogene (melanoma growth-stimulating activity)|MGSA alpha|chemokine (C-X-C motif) ligand 1|fibroblast secretory protein|melanoma growth stimulatory activity alpha

Pathway

- Chemokine signaling pathway
- Cytokine-cytokine receptor interaction
- Epithelial cell signaling in Helicobacter pylori infection

Disease

- Alzheimer disease
- Asthma
- Bronchiolitis
- Genetic Predisposition to Disease
- Infant

Copyright © 2023 Abnova Corporation. All Rights Reserved.

- Ovarian Neoplasms
- Respiratory Syncytial Virus Infections

Copyright © 2023 Abnova Corporation. All Rights Reserved.