
ATF2 (phospho T71) monoclonal antibody, clone FOO-1

Catalog # MAB20331 Size 100 uL

Applications



Western Blot (Cell lysate)

Western blot analysis of ATF2 (phospho T71) expression in (1) HeLa cell lysate; (2) HeLa cell lysate treated with Anisomycin.

Specification

Product Description	Rabbit monoclonal antibody raised against synthetic phosphopeptide of human ATF2.
Immunogen	A synthetic phosphopeptide corresponding to residues surrounding T71 of human ATF2.
Host	Rabbit
Reactivity	Human
Form	Liquid
Purification	Affinity purification
Isotype	IgG
Recommend Usage	Immunocytochemistry (1:50-1:200) Immunofluorescence (1:50-1:200) Immunoprecipitation (1:30) Western Blot (1:500-1:2000) The optimal working dilution should be determined by the end user.
Storage Buffer	In PBS, 150 mM NaCl, pH 7.4 (50% glycerol, 0.4-0.5 mg/mL BSA, 0.02% sodium azide).

Copyright © 2023 Abnova Corporation. All Rights Reserved.

Storage Instruction	Store at -20°C for one year. After reconstitution, at 4°C for one month. It can also be aliquotted and st ored frozen at -20°C for a longer time. Avoid repeated freezing and thawing.
Note	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which shoul d be handled by trained staff only.

Applications

- Immunofluorescence
- Immunocytochemistry
- Immunoprecipitation
- Western Blot (Cell lysate)

  Western blot analysis of ATF2 (phospho T71) expression in (1) HeLa cell lysate; (2) HeLa cell lysate treated with Anisomycin.

Gene Info — ATF2

Entrez GeneID	1386
Protein Accession#	P15336
Gene Name	ATF2
Gene Alias	CRE-BP1, CREB2, HB16, MGC111558, TREB7
Gene Description	activating transcription factor 2
Omim ID	123811
Gene Ontology	Hyperlink
Gene Summary	This gene encodes a transcription factor that is a member of the leucine zipper family of DNA bin ding proteins. This protein binds to the cAMP-responsive element (CRE), an octameric palindrom e. The protein forms a homodimer or heterodimer with c-Jun and stimulates CRE-dependent tran scription. The protein is also a histone acetyltransferase (HAT) that specifically acetylates histone s H2B and H4 in vitro; thus it may represent a class of sequence-specific factors that activate tran scription by direct effects on chromatin components. Additional transcript variants have been iden tified but their biological validity has not been determined. [provided by RefSeq
Other Designations	OTTHUMP00000163262|activating transcription factor 2 splice variant ATF2-var2|cAMP respons ive element binding protein 2, formerly

Copyright © 2023 Abnova Corporation. All Rights Reserved.

Pathway

- MAPK signaling pathway

Disease

- Bipolar Disorder
- Cardiovascular Diseases
- Diabetes Mellitus
- Edema
- Genetic Predisposition to Disease
- Schizophrenia
- Tobacco Use Disorder

Copyright © 2023 Abnova Corporation. All Rights Reserved.