
LATS1/LATS2 (phospho T1079/1041) polyclonal antibody

Catalog # PAB29206 Size 100 uL

Applications



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical staining of human brain tissue by LATS1/LATS2 (phospho T1079/1041) polyclonal antibody (Cat # PAB29206) without blocking peptide (A) or preincubated with blocking peptide (B) under 1:50-1:100 dilution.

Specification

Product Description	Rabbit polyclonal antibody raised against synthetic phosphopeptide of human LATS1/LATS2.
Immunogen	Synthetic phosphopeptide (conjugated with KLH) corresponding to residues surrounding T1079/1041 of human LATS1/LATS2.
Host	Rabbit
Theoretical MW (kDa)	126
Reactivity	Human, Mouse
Specificity	This antibody detects endogenous levels of LATS1/LATS2 only when phosphorylated at threonine 1079/1041.
Form	Liquid

Copyright © 2023 Abnova Corporation. All Rights Reserved.

Purification	Affinity purification
Recommend Usage	Immunohistochemistry (1:50-1:100) The optimal working dilution should be determined by the end user.
Storage Buffer	In PBS, pH 7.4 (without Mg^{2+} and Ca^{2+}), (50% glycerol, 0.02% sodium azide)
Storage Instruction	Store at -20°C. Avoid repeated freezing and thawing.

Applications

- Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

  Immunohistochemical staining of human brain tissue by LATS1/LATS2 (phospho T1079/1041) polyclonal antibody (Cat # PAB29206) without blocking peptide (A) or preincubated with blocking peptide (B) under 1:50-1:100 dilution.

Gene Info — LATS1

Entrez GeneID	9113
Protein Accession#	O95835;Q9NRM7
Gene Name	LATS1
Gene Alias	WARTS, wts
Gene Description	LATS, large tumor suppressor, homolog 1 (Drosophila)
Omim ID	603473
Gene Ontology	Hyperlink
Gene Summary	The protein encoded by this gene is a putative serine/threonine kinase that localizes to the mitotic apparatus and complexes with cell cycle controller CDC2 kinase in early mitosis. The protein is p hosphorylated in a cell-cycle dependent manner, with late prophase phosphorylation remaining thr ough metaphase. The N-terminal region of the protein binds CDC2 to form a complex showing re duced H1 histone kinase activity, indicating a role as a negative regulator of CDC2/cyclin A. In ad dition, the C-terminal kinase domain binds to its own N-terminal region, suggesting potential nega tive regulation through interference with complex formation via intramolecular binding. Biochemica l and genetic data suggest a role as a tumor suppressor. This is supported by studies in knockout mice showing development of soft-tissue sarcomas, ovarian stromal cell tumors and a high sensiti vity to carcinogenic treatments. [provided by RefSeq
Other Designations	LATS (large tumor suppressor, Drosophila) homolog 1|LATS homolog 1

Copyright © 2023 Abnova Corporation. All Rights Reserved.

Gene Info — LATS2

Entrez GeneID	26524
Protein Accession#	O95835;Q9NRM7
Gene Name	LATS2
Gene Alias	FLJ13161, KPM
Gene Description	LATS, large tumor suppressor, homolog 2 (Drosophila)
Omim ID	604861
Gene Ontology	Hyperlink
Gene Summary	This gene encodes a serine/threonine protein kinase belonging to the LATS tumor suppressor fa mily. The protein localizes to centrosomes during interphase, and early and late metaphase. It inte racts with the centrosomal proteins aurora-A and ajuba and is required for accumulation of gamm a-tubulin and spindle formation at the onset of mitosis. It also interacts with a negative regulator of p53 and may function in a positive feedback loop with p53 that responds to cytoskeleton damage. Additionally, it can function as a co-repressor of androgen-responsive gene expression. [provided by RefSeq
Other Designations	LATS (large tumor suppressor, Drosophila) homolog 2|LATS, large tumor suppressor, homolog 2| OTTHUMP00000018106|kinase phosphorylated during mitosis protein|serine/threonine kinase K PM|warts-like kinase

Disease

- Adenocarcinoma
- Esophageal Neoplasms
- Tobacco Use Disorder

Copyright © 2023 Abnova Corporation. All Rights Reserved.