
RecomAb™

HDAC1 recombinant monoclonal antibody, clone 10A1

Catalog # RAB04027 Size 100 uL

Applications



Western Blot

Western Blot analysis of Lane 1: Hela whole cell lysate; Lane 2: K562 whole cell lysate; Lane 3: Jurkat whole cell lysate; Lane 4: HL60 whole cell lysate.

Specification

Product Description	Rabbit recombinant monoclonal antibody raised against human HDAC1.
Antibody Species	Rabbit
Immunogen	Original antibody is raised against recombinant protein corresponding to full length human HDAC1.
Reactivity	Human
Form	Liquid
Preparation Method	Mammalian cell (HEK293) expression system
Purification	Affinity-chromatography
Isotype	IgG

Copyright © 2023 Abnova Corporation. All Rights Reserved.

Recommend Usage	ELISA Western Blot (1:500-1:5000) The optimal working dilution should be determined by the end user.
Storage Buffer	In PBS, pH7.4 (150mM NaCl, 50% glycerol and 0.02% sodium azide)
Storage Instruction	store at -20 °C or -80 °C. Aliquot to avoid repeated freezing and thawing.
Note	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which shoul d be handled by trained staff only.

Applications

- Enzyme-linked Immunoabsorbent Assay
- Western Blot

  Western Blot analysis of Lane 1: Hela whole cell lysate; Lane 2: K562 whole cell lysate; Lane 3: Jurkat whole cell lysate; Lane 4: HL60 whole cell lysate.

Gene Info — HDAC1

Entrez GeneID	3065
Protein Accession#	Q13547
Gene Name	HDAC1
Gene Alias	DKFZp686H12203, GON-10, HD1, RPD3, RPD3L1
Gene Description	histone deacetylase 1
Omim ID	601241
Gene Ontology	Hyperlink
Gene Summary	Histone acetylation and deacetylation, catalyzed by multisubunit complexes, play a key role in the regulation of eukaryotic gene expression. The protein encoded by this gene belongs to the histon e deacetylase/acuc/apha family and is a component of the histone deacetylase complex. It also in teracts with retinoblastoma tumor-suppressor protein and this complex is a key element in the con trol of cell proliferation and differentiation. Together with metastasis-associated protein-2, it deac etylates p53 and modulates its effect on cell growth and apoptosis. [provided by RefSeq
Other Designations	OTTHUMP00000008745|reduced potassium dependency, yeast homolog-like 1

Copyright © 2023 Abnova Corporation. All Rights Reserved.

Pathway

- Cell cycle
- Chronic myeloid leukemia
- Notch signaling pathway
- Pathways in cancer

Disease

- Asthma
- Cognition Disorders
- Genetic Predisposition to Disease
- Huntington disease
- Mental Status Schedule
- Neoplasms
- Ovarian cancer
- Ovarian Neoplasms
- Tobacco Use Disorder

Copyright © 2023 Abnova Corporation. All Rights Reserved.