
RecomAb™

ACVR2B recombinant monoclonal antibody, clone 7B10

Catalog # RAB07543 Size 100 uL

Applications



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical analysis of paraffin-embedded human breast cancer with ACVR2B recombinant monoclonal antibody, clone 7B10 (Cat # RAB07543) on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a Goat anti-rabbit IgG polymer labeled by HRP and visualized using 0.05% DAB.

Specification

Product Description	Rabbit recombinant monoclonal antibody raised against human ACVR2B.
Antibody Species	Rabbit
Immunogen	Original antibody is raised against a synthetic peptide corresponding to human ACVR2B.
Reactivity	Human
Form	Liquid
Preparation Method	Mammalian cell (HEK293) expression system
Purification	Affinity chromatography purification
Isotype	IgG
Recommend Usage	ELISA Immunohistochemistry (1:50-1:200) The optimal working dilution should be determined by the end user.

Copyright © 2023 Abnova Corporation. All Rights Reserved.

Storage Buffer	In PBS, pH7.4 (150 mM NaCl, 0.02% sodium azide and 50% glycerol)
Storage Instruction	Store at -20°C or -80°C. Aliquot to avoid repeated freezing and thawing.
Note	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which shoul d be handled by trained staff only.

Applications

- Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

  Immunohistochemical analysis of paraffin-embedded human breast cancer with ACVR2B recombinant monoclonal antibody, clone 7B10 (Cat # RAB07543) on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a Goat anti-rabbit IgG polymer labeled by HRP and visualized using 0.05% DAB.

- Enzyme-linked Immunoabsorbent Assay

Gene Info — ACVR2B

Entrez GeneID	93
Protein Accession#	Q13705
Gene Name	ACVR2B
Gene Alias	ACTRIIB, ActR-IIB, MGC116908
Gene Description	activin A receptor, type IIB
Omim ID	602730
Gene Ontology	Hyperlink
Gene Summary	Activins are dimeric growth and differentiation factors which belong to the transforming growth fac tor-beta (TGF-beta) superfamily of structurally related signaling proteins. Activins signal through a heteromeric complex of receptor serine kinases which include at least two type I (I and IB) and two type II (II and IIB) receptors. These receptors are all transmembrane proteins, composed of a ligan d-binding extracellular domain with cysteine-rich region, a transmembrane domain, and a cytopla smic domain with predicted serine/threonine specificity. Type I receptors are essential for signalin g; and type II receptors are required for binding ligands and for expression of type I receptors. Typ e I and II receptors form a stable complex after ligand binding, resulting in phosphorylation of type I receptors by type II receptors. Type II receptors are considered to be constitutively active kinases. This gene encodes activin A type IIB receptor, which displays a 3- to 4-fold higher affinity for the li gand than activin A type II receptor. [provided by RefSeq

Copyright © 2023 Abnova Corporation. All Rights Reserved.

Other Designations	activin A type IIB receptor

Pathway

- Cytokine-cytokine receptor interaction
- TGF-beta signaling pathway

Disease

- Genetic Predisposition to Disease
- Hyperparathyroidism
- Obesity
- Ovarian Failure
- Polycystic Ovary Syndrome
- Puberty
- Thrombophilia
- Tobacco Use Disorder

Copyright © 2023 Abnova Corporation. All Rights Reserved.