
MaxPab®

ACVR2B purified MaxPab mouse polyclonal antibody (B01P)

Catalog # H00000093-B01P Size 50 ug

Applications



Western Blot (Transfected lysate)

Western Blot analysis of ACVR2B expression in transfected 293T cell line (H00000093-T02) by ACVR2B MaxPab polyclonal antibody.

Lane 1: ACVR2B transfected lysate(34.20 KDa).
Lane 2: Non-transfected lysate.

Specification

Product Description	Mouse polyclonal antibody raised against a full-length human ACVR2B protein.
Immunogen	ACVR2B (AAH96245.1, 1 a.a. ~ 303 a.a) full-length human protein.
Sequence	MNDFVAVKIFPLQDKQSWQSEREIFSTPGMKHENLLQFIAAEKRGSNLEVELWLITAFHDKGSLTD YLKGNIITWNELCHVAETMSRGLSYLHEDVPWCRGEGHKPSIAHRDFKSKNVLLKSDLTAVLADF GLAVRFEPGKPPGDTHGQVGTRRYMAPEVLEGAINFQRDAFLRIDMYAMGLVLWELVSRCKAAD GPVDEYMLPFEEEIGQHPSLEELQEVVVHKKMRPTIKDHWLKHPGLAQLCVTIEECWDHDAEAR LSAGCVEERVSLIRRSVNGTTSDCLVSLVTSVTNVDLPPKESSI
Host	Mouse
Reactivity	Human
Interspecies Antigen Sequence	Mouse (99); Rat (99)
Quality Control Testing	Antibody reactive against mammalian transfected lysate.
Storage Buffer	In 1x PBS, pH 7.4
Storage Instruction	Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Copyright © 2023 Abnova Corporation. All Rights Reserved.

Applications

- Western Blot (Transfected lysate)

  Western Blot analysis of ACVR2B expression in transfected 293T cell line (H00000093-T02) by ACVR2B MaxPab polyclonal antibody.

  Lane 1: ACVR2B transfected lysate(34.20 KDa).
  Lane 2: Non-transfected lysate.

  Protocol Download

Gene Info — ACVR2B

Entrez GeneID	93
GeneBank Accession#	BC096245.1
Protein Accession#	AAH96245.1
Gene Name	ACVR2B
Gene Alias	ACTRIIB, ActR-IIB, MGC116908
Gene Description	activin A receptor, type IIB
Omim ID	602730
Gene Ontology	Hyperlink
Gene Summary	Activins are dimeric growth and differentiation factors which belong to the transforming growth fac tor-beta (TGF-beta) superfamily of structurally related signaling proteins. Activins signal through a heteromeric complex of receptor serine kinases which include at least two type I (I and IB) and two type II (II and IIB) receptors. These receptors are all transmembrane proteins, composed of a ligan d-binding extracellular domain with cysteine-rich region, a transmembrane domain, and a cytopla smic domain with predicted serine/threonine specificity. Type I receptors are essential for signalin g; and type II receptors are required for binding ligands and for expression of type I receptors. Typ e I and II receptors form a stable complex after ligand binding, resulting in phosphorylation of type I receptors by type II receptors. Type II receptors are considered to be constitutively active kinases. This gene encodes activin A type IIB receptor, which displays a 3- to 4-fold higher affinity for the li gand than activin A type II receptor. [provided by RefSeq
Other Designations	activin A type IIB receptor

Pathway

Copyright © 2023 Abnova Corporation. All Rights Reserved.

- Cytokine-cytokine receptor interaction
- TGF-beta signaling pathway

Disease

- Genetic Predisposition to Disease
- Hyperparathyroidism
- Obesity
- Ovarian Failure
- Polycystic Ovary Syndrome
- Puberty
- Thrombophilia
- Tobacco Use Disorder

Copyright © 2023 Abnova Corporation. All Rights Reserved.