RecomAb™ Sm Size

Histone H3 (phospho T11) monoclonal antibody, clone RM164

Catalog # RAB09462 Size 25 ug

Applications



Western Blot (Cell lysate)

Western Blot analysis of Lane 1: acid extracts of HeLa cell treated with Nocodazole and Lane 2: acid extracts of HeLa cell untreated with Histone H3 (phospho T11) recombinant monoclonal antibody, clone RM164 (Cat # RAB09462) at 0.5 ug/mL working concentration, showed a band of Histone H3 phosphorylated at Threonine 11.

Specification

Product Description	Rabbit recombinant monoclonal antibody raised against of human histone H3 (T11).
Antibody Species	Rabbit
Immunogen	Original antibody is raised against a synthetic phosphopeptide corresponding to residues surroundin g T11 of human Histone H3.
Sequence	N/A
Specificity	This antibody reacts to Histone H3 phosphorylated at Threonine 11. No cross reactivity with other ph osphorylated histones.
Form	Liquid
Purification	Protein A purification
Isotype	IgG
Recommend Usage	ELISA (0.2 ug/mL-1 ug/mL) Western Blot (0.5 ug/mL-2 ug/mL) The optimal working dilution should be determined by the end user.

Copyright © 2023 Abnova Corporation. All Rights Reserved.

Storage Buffer	In PBS (50% glycerol, 1% BSA, 0.09% sodium azide)
Storage Instruction	Store at -20°C. Aliquot to avoid repeated freezing and thawing.
Note	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which shoul d be handled by trained staff only.

Applications

- Enzyme-linked Immunoabsorbent Assay
- Western Blot (Cell lysate)

  Western Blot analysis of Lane 1: acid extracts of HeLa cell treated with Nocodazole and Lane 2: acid extracts of HeLa cell untreated with Histone H3 (phospho T11) recombinant monoclonal antibody, clone RM164 (Cat # RAB09462) at 0.5 ug/mL working concentration, showed a band of Histone H3 phosphorylated at Threonine 11.

Gene Info — HIST1H3A

Entrez GeneID	8350
Protein Accession#	P84243
Gene Name	HIST1H3A
Gene Alias	H3/A, H3FA
Gene Description	histone cluster 1, H3a
Omim ID	602810
Gene Ontology	Hyperlink
Gene Summary	Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chro mosomal fiber in eukaryotes. This structure consists of approximately 146 bp of DNA wrapped ar ound a nucleosome, an octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H 1, with the DNA between the nucleosomes to form higher order chromatin structures. This gene is intronless and encodes a member of the histone H3 family. Transcripts from this gene lack polyA t ails; instead, they contain a palindromic termination element. This gene is found in the large histon e gene cluster on chromosome 6p22-p21.3. [provided by RefSeq
Other Designations	H3 histone family, member A|histone 1, H3a

Copyright © 2023 Abnova Corporation. All Rights Reserved.

Pathway

- Systemic lupus erythematosus

Copyright © 2023 Abnova Corporation. All Rights Reserved.