Bioactive

PPBP (Human) Recombinant Protein

Catalog # P9554 Size 2 x 10 ug

Specification

Product Description	Human PPBP (P02775, 59 a.a. - 128 a.a.) partial recombinant protein expressed in *Escherichia coli* .
Sequence	AELRCMCIKTTSGIHPKNIQSLEVIGKGTHCNQVEVIATLKDGRKICLDPDAPRIKKIVQKKLAGDES AD
Host	Escherichia coli
Theoretical MW (kDa)	7.6
Form	Lyophilized
Preparation Method	*Escherichia coli* expression system
Purity	> 97.0% by SDS-PAGE
Activity	The Biological activity is 1 - 10 ng/mL, was determined by the ability of NAP2 to chemoattract human neurotrophils, corresponding to a specific activity of 100000 - 1000000 IU/mg.
Recommend Usage	Biological Activity SDS-PAGE The optimal working dilution should be determined by the end user.
Storage Buffer	Lyophilized from sterile distilled Water is > 100 ug/mL
Storage Instruction	Store at 2°C to 8°C for 1 week. For long term storage, aliquot and store at -20°C to -80°C. Aliquot to avoid repeated freezing and thawing.

Applications

- Functional Study
- SDS-PAGE

Copyright © 2023 Abnova Corporation. All Rights Reserved.

Gene Info — PPBP

Entrez GeneID	5473
Protein Accession#	P02775
Gene Name	PPBP
Gene Alias	B-TG1, Beta-TG, CTAP-III, CTAP3, CTAPIII, CXCL7, LA-PF4, LDGF, MDGF, NAP-2, PBP, SCYB7, TC1, TC2, TGB, TGB1, THBGB, THBGB1
Gene Description	pro-platelet basic protein (chemokine (C-X-C motif) ligand 7)
Omim ID	121010
Gene Ontology	Hyperlink
Gene Summary	The protein encoded by this gene is a platelet-derived growth factor that belongs to the CXC chemokine family. This growth factor is a potent chemoattractant and activator of neutrophils. It has been shown to stimulate various cellular processes including DNA synthesis, mitosis, glycolysis, intracellular cAMP accumulation, prostaglandin E2 secretion, and sythesis of hyaluronic acid and sulfated glycosaminoglycan. It also stimulates the formation and secretion of plasminogen activator by synovial cells. [provided by RefSeq
Other Designations	CXC chemokine ligand 7|beta-thromboglobulin|connective tissue-activating peptide III|leukocyte-derived growth factor|low-affinity platelet factor IV|macrophage-derived growth factor|neutrophil-activating peptide 2|neutrophil-activating peptide-2|platelet

Pathway

- Chemokine signaling pathway
- Cytokine-cytokine receptor interaction

Disease

- Asthma
- Bronchiolitis
- Cardiovascular Diseases
- Diabetes Mellitus
- Edema

Copyright © 2023 Abnova Corporation. All Rights Reserved.

- Genetic Predisposition to Disease
- Infant

- Respiratory Syncytial Virus Infections

Copyright © 2023 Abnova Corporation. All Rights Reserved.